







Dignity Health Quality Transparency Dashboard

Saint Francis Memorial Hospital

At Dignity Health Saint Francis Memorial Hospital, our mission is to provide high quality compassionate health care for all who seek it, and we strive for the best, safest care for our patients. We're participating in the Hospital Quality Transparency Dashboard project to enable consumers to make informed choices about their health care through publicly available data about our hospital, and we encourage others to do the same. Below are our recent results for five outcome measures out of hundreds that we track, showing how we compare to other hospitals in California and across the U.S. **Remember, lower numbers indicate better patient care for that specific measure.**

OUTCOME MEASURES

 Lower is better	 CLABSI	 Colon SSI	 Low-risk Cesarean Rate	 VTE	 Sepsis Mortality
Saint Francis Memorial Hospital	.73	N/A	N/A	0.00%	11.6%
Measure Period	1/01/2018 – 6/30/2018				1/01/2017 - 12/31/2017
California State Level	.85	.97	13.3%	3.0%	14.9%
Measure Period	1/01/2017 - 12/31/2017				
National Level	.89	.93	12.7%	2.0%	25.0%
Measure Period	1/01/2016 - 12/31/2016		1/01/2011 - 12/31/2011	1/01/2017 - 12/31/2017	

PROGRAM STATUS

<input type="checkbox"/> Yes <input checked="" type="checkbox"/> Deliveries are not available at this hospital	Maternity Safety Program (Program Status Measure) This hospital has a Maternity Safety Program in place. A Maternity Safety Program provides a coordinated approach and emergency response to risks associated with pregnancy and childbirth.
<input checked="" type="checkbox"/> Yes	Sepsis Protocol (Program Status Measure) This hospital has a Sepsis Protocol in place. This offers a protocolized approach to identifying and treating patients with severe sepsis (infection with a systemic inflammatory response, the body's response to an infection, and an acute organ dysfunction, organ does not perform expected function) or septic shock (persistent sepsis induced tissue hypoperfusion or dangerously low blood flow through an organ) in the emergency department or acute care in-patient setting.
<input checked="" type="checkbox"/> Yes	Respiratory Monitoring Program (Program Status Measure) This hospital has a Respiratory Monitoring Program in place. Respiratory monitoring provides guidance for assessment of risk of respiratory depression, and slow and ineffective breathing, and it includes continuous monitoring of breathing and functioning of the lungs and circulatory system when indicated.

Dignity Health Quality Transparency Dashboard

OUTCOME MEASURE DEFINITIONS



CLABSI – CENTRAL LINE-ASSOCIATED BLOOD STREAM INFECTION

A central line-associated blood stream infection is a serious infection that occurs when bacteria or germs enter the bloodstream through an IV called a central line. A central line allows access to a major vein close to the heart and can stay in place for weeks or months. It is mainly used for medication delivery and blood draws. Central line infections are measured using the Centers for Disease Control and Prevention Standardized Infection Ratio (SIR). This ratio is the observed number of infections divided by the number of expected infections. Values less than 1.00 show that the number of observed infections was lower than expected. Values more than 1.00 show that the number of infections was higher than expected. N/A means the scoring related to predicted infections is not available.



COLON SSI – SURGICAL SITE INFECTION FROM COLON SURGERY

A surgical site infection occurs after surgery and is caused by bacteria or germs in the part of the body where the surgery took place. Surgical site infections for colon surgery are measured using the Centers for Disease Control and Prevention Standardized Infection Ratio (SIR). This ratio is the observed number of infections divided by the number of expected infections. Values less than 1.00 show that the number of observed infections was lower than expected. Values more than 1.00 show that the number of infections was higher than expected. N/A means the scoring related to predicted infections is not available.



LOW-RISK CESAREAN BIRTHS

Safely reducing cesarean deliveries in low-risk pregnancies is a priority for hospitals. The low-risk cesarean birth rate is the number of cesarean or surgical births divided by the number of all births. This measurement includes all moms who deliver without complicated or “high risk pregnancies.” It excludes patients with twin pregnancies, preterm delivery, previous cesarean, and other high-risk conditions that often require a cesarean. A lower number indicates fewer cesareans were performed among low-risk mothers.



VTE – Preventable Venous Thromboembolisms

Staying in bed for long periods of time can put hospital patients at an increased risk for developing a blood clot known as a venous thromboembolism, or VTE. There are safe, effective, and proven procedures to prevent blood clots or to treat them when they occur. Patients at risk for VTEs should receive these preventable measures during their hospital stay. Preventable VTEs are measured by the percent of patients who developed a blood clot while in the hospital and did not receive treatment that could have prevented it. A lower number shows a better outcome.



SEPSIS MORTALITY

Sepsis or septic shock is the body’s extreme response to an infection. It is a life-threatening medical emergency that is a risk for patients admitted to an emergency department or hospital care setting. Without timely treatment, sepsis can lead to tissue damage, organ failure (e.g., kidney, lungs, liver, etc.), and death. Sepsis infection is measured by the percent of patients who die in the hospital. A lower number shows a better outcome.