

Anesthesia Preoperative Testing Guidelines

Patient Category	СВС	Chem 8	Chem 14	PT/INR	PTT	UA	Urine HCG	ECG	CXR
Age over 50 years								Х	
Cardiac Disease (MI, CHF, Pacemaker/AICD, Coronary Stents)	x	Х						Х	
Pulmonary Disease (COPD)	X							Х	X
Diabetes	Х	Х						Х	
Hepatic Disease	X		Х	Х	Х				
Hypertension	X	Х						X	
Morbid (Extreme) Obesity	X	Х		Х				X	
Neurological Disease	X	Х							
ESRD on Hemodialysis	X	Х		Х	Х			Х	
Renal Insufficiency	X	Х							
Vascular Disease	X							Х	
UTI						Х			
Anticoagulant (Coumadin)	X			Х					
Anticoagulant (Heparin)					X				
Chemotherapy	X								
Diuretics		Х							
Cardiac/Thoracic Surgery	X		Х	Х	Х			Х	X
Major Surgery (ref. PC-1860)	X	Х		Х				Х	X
Menstruating Female < 50 yrs							Х		

- Patients with ESRD on Hemodialysis must have Chem 8 within 24 hours of the scheduled procedure.
- All patients with diabetes must have glucose (serum or FSBS) done on the day of surgery.
- * All patients on Coumadin must have PT/INR done on the day of the procedure.
- * All patients on Heparin must have PTT done on the day of the procedure.
- ✤ All menstruating females must have HCG done on the day of the scheduled procedure.
- ECG and CXR will be accepted within 1 year unless there are changes in cardiac or respiratory symptoms.
- * All other labs must be completed within 30 days of the scheduled procedure.
- All abnormal labs are to be reviewed and addressed by the surgeon prior to the day of surgery.
- ♦ History and Physical must be completed within 30 days of the procedure (ref. MS-30).

Based on patient assessment, more tests may be required.

These guidelines are approved by Department of Anesthesia (for patients receiving procedural sedation, tests will be ordered by the interventionalist).